



Listing of Useful Plants of the World


[formerly Listing of Potential New Crops for Australia]

NOTICE: Information on the [background to this Listing](#) is available. The analysis of numbers of papers/mentions over time was completed in 1997, using the Agricola database (1970-1996). Hard copies of the Listing (553 pages; over 4200 new crops listed with the analyses of numbers of papers over time included) are available from the publisher, Dr Rob Fletcher; see [Advice on Publications Available](#).

The references listed below are from the Biological Abstracts database (1988-2000) and are courtesy of SilverPlatter Information. For more information re Silverplatter, go to www.silverplatter.com.

Ficus carica

Analysis of numbers of papers/mentions over time (Agricola database 1970-1996):



Source: AGRICOLA database (1970-1996)

Common Name(s): fig syn figure syn higo syn feige

Crop Use(s): fruit

Reference Source(s): rehm, sturtevant

Number of Papers/Mentions: 175

References (Biological Abstracts 1988-2000):

[These references are from the Biological Abstracts database and are courtesy of SilverPlatter Information. For more information re SilverPlatter, go to www.silverplatter.com. References are listed alphabetically by author within years, with most recent references first; addresses of author(s) have been included when available. To search within this page, we suggest using Find in Page, within the Edit menu of the Web Browser.]

Canal, J. R., M. D. Torres, et al. (2000). A chloroform extract obtained from a decoction of *Ficus carica* leaves improves the cholesterolaemic status of rats with streptozotocin-induced diabetes. *Acta Physiologica Hungarica*. [print] 87(1): 71-76. {a} Departamento de Fisiologia, Facultad de Medicina, Universidad de Extremadura, Avenida de Elvas s/n, 06007, Badajoz, Spain

Kang, H., M. Y. Kang, et al. (2000). Identification of natural rubber and characterization of rubber biosynthetic activity in fig tree. *Plant Physiology Rockville*. [print] July 123(3): 1133-1142. {a} Kumho Life and Environmental Science Laboratory, 1 Oryong-dong, Puk-gu, Kwangju, 500-712, South Korea

Kanzaki, N., K. Tsuda, et al. (2000). Description of *Bursaphelenchus conicaudatus* n. sp. (Nematoda: Aphelenchoididae), isolated from the yellow-spotted longicorn beetle, *Psacotha hilaris* (Coleoptera: Cerambycidae) and fig trees, *Ficus carica*. *Nematology* . [print] 2(2): 165-168. {a} Graduate School of

Agriculture, Kyoto University, Sakyo-ku, Kyoto, 606-8502, Japan

Perez, C., E. Dominguez, et al. (2000). Hypoglycaemic activity of an aqueous extract from *Ficus carica* (Fig Tree) leaves in streptozotocin diabetic rats. *Pharmaceutical Biology*. [print] July 38(3): 181-186. {a} Departamento de Fisiologia, Facultad de Medicina, Universidad de Extremadura, Avenida de Elvas, s/n, 06071, Badajoz, Spain

Teragishi, A., Y. Kanbara, et al. (2000). Effects of foliar application of choline chloride on the quality of winter-cropped fig cv. Masui-Dauphine grown in hydroponics. *Journal of the Japanese Society for Horticultural Science*. [print] July 69(4): 390-395. {a} Kyoto Prefectural Agricultural Promotion and Distribution Division, Kamigyo-ku, Kyoto, 602-8570, Japan

Wenninger, P. S. and L. A. Shipley (2000). Harvesting, rumination, digestion, and passage of fruit and leaf diets by a small ruminant, the blue duiker. *Oecologia Berlin*. [print] June 123(4): 466-474. {a} Department of Natural Resource Sciences, Washington State University, Pullman, WA, 99164-6410, USA

Yahata, D. and H. Nogata (2000). Changes in sugar and polyuronide contents in fig syconia during storage. *Nippon Shokuhin Kagaku Kogaku Kaishi*. [print] 47(4): 333-337. {a} Buzen Branch, Fukuoka Agricultural Research Center, 2-4-1, Nishiizumi, Yukuhashi, Fukuoka, 824-0038, Japan

Yahata, D. and H. Nogata (2000). Relationship between air temperature and characteristics of developing fig syconia at different nodal positions. *Journal of the Japanese Society for Horticultural Science*. March 69(2): 202-207. {a} Buzen Branch, Fukuoka Agricultural Research Center, Nishiizumi, Yukuhashi, Fukuoka, 824-0038, Japan

Batanouny, K. H. (1999). The Mediterranean coastal dunes in Egypt: An endangered landscape. *Estuarine Coastal and Shelf Science*. Aug. 49(Suppl. A): 3-9. {a} Faculty of Science, University of Cairo, Giza, Egypt

Cowart, N. M. and J. H. Graham (1999). Within- and among-individual variation in fluctuating asymmetry of leaves in the fig (*Ficus carica* L.). *International Journal of Plant Sciences* 160(1): 116-121. {a} Department of Biology, Berry College, Mount Berry, GA, 30149-0446, USA

de, A. A., H. R. Borba, et al. (1999). Anthelmintic activity of the latex of *Ficus* species. *Journal of Ethnopharmacology*. March 64(3): 255-258. {a} Empresa Brasileira de Pesquisa Agropecuaria, Centro Nacional de Pesquisa de Tecnologia Agroindustrial de Alimentos, Av. das Americas 29501, 23020-470, Rio de Janeiro, Brazil

Galderisi, U., M. Cipollaro, et al. (1999). Identification of the edible fig 'Bianco del Cilento' by random amplified polymorphic DNA analysis. *Hortscience*. Dec. 34(7): 1263-1265. {a} Centro di Ricerca Interdipartimentale di Scienze Computazionali e Biotecnologiche (C.R.I.S.C.E.B.), Facolta di Medicina e Chirurgia, Istituto di Farmacologia e Tossicologia, 2.a Universita degli Studi di Napoli, Via Costantinopoli 16, 80138, Naples, Italy

- Mendez, E. (1999). Fruit trees in the streets: Mendoza, Argentina. *Revista de la Facultad de Ciencias Agrarias Universidad Nacional de Cuyo* 31(2): 21-30. {a} UID Botanica y Fitosociologia, IADIZA-CRICYT, CONICET, Av. Adrian Ruiz Leal s/n, Parque Gral, San Martin, Mendoza, 5500, Argentina
- Neilson, R. and D. J. F. Brown (1999). Feeding on different host plants alters the natural abundances of delta13C and delta15N in Longidoridae (Nemata). *Journal of Nematology*. March 31(1): 20-26. {a} Nematology Unit, Scottish Crop Research Institute, Dundee, DD2 5DA, UK
- Nogales, M., E. C. Hernandez, et al. (1999). Seed dispersal by common ravens *Corvus corax* among island habitats (Canarian Archipelago). *Ecoscience* 6(1): 56-61. {a} Departamento de Biologia Animal (Zoologia), 38206, Universidad de La Laguna, Tenerife, Canary Islands, Spain
- Pareek, O. P. (1999). Arid zone fruits research in India. *Indian Journal of Agricultural Sciences*. Aug. 68 (8 spec. issue): 508-514. {a} National Research Centre for Arid Horticulture, Bikaner, RAJ, 334 003, India
- Stika, H. P. (1999). Approaches to reconstruction of early Celtic land-use in the central Neckar region in southwestern Germany. *Vegetation History and Archaeobotany*. June 8(1-2): 95-103. {a} Institut fuer Botanik (210), Universitaet Hohenheim, D-70593, Stuttgart, Germany
- Vinson, J. A. (1999). The functional food properties of figs. *Cereal Foods World* 44(2): 82-87. {a} Univ. Scranton, Scranton, PA, USA
- Wang, L. J., Y. L. Liu, et al. (1999). The role of calcium in promotion of proline accumulation induced by salt stress in fig (*Ficus carica* L.) cells. *Acta Phytophysiological Sinica* 25(1): 38-42. {a} Department of Agronomy, Nanjing Agricultural University, Nanjing, 210095, China
- Yahata, D. and H. Nogata (1999). Cultivar variations in sugar contents in the fig *syconia*, their parts and nodal positions. *Journal of the Japanese Society for Horticultural Science*. Sept. 68(5): 987-992. {a} Buzen Branch, Fukuoka Agricultural Research Center, Nishiizumi, Yukuhashi, Fukuoka, 824-0038, Japan
- Atzmon, N. and Z. Henkin (1998). Establishing forest tree species on peatland in a reflooded area of the Huleh Valley, Israel. *Forestry Oxford* 71(2): 141-146. {a} Inst. Field Garden Crops, Agric. Res. Organization, Volcani Cent., P.O. Box 6, Bet Dagan 50250, Israel
- Brehler, R., E. Abrams, et al. (1998). Cross-reactivity between *Ficus benjamina* (weeping fig) and natural rubber latex. *Allergy Copenhagen* 53(4): 402-406. {a} Dep. of Dermatology, Von-Esmarch-Str. 56, 48149 Muenster, Germany
- Doster, M. A. and T. J. Michailides (1998). Production of bright greenish yellow fluorescence in figs infected by *Aspergillus* species in California orchards. *Plant Disease* 82(6): 669-673. Dep. Plant Pathol., Univ. Calif., Davis, CA, USA

- Gibernau, M., M. M. Hossaert, et al. (1998). Are olfactory signals sufficient to attract fig pollinators? *Ecoscience* 5(3): 306-311. {a} CEFE/CNRS, 1919 Route de Mende, 34293 Montpellier Cedex 5, France
- Gonzalez, V., J. Ramallo, et al. (1998). Fig fruit rot caused by *Phytophthora palmivora* in fig crops in Tucuman, Argentina. *Fitopatologia* 33(4): 228-231. {a} Estacion Experimental Agroindustrial Obispo Colombres, 4101 Las Talitas, Tucuman, Argentina
- Hosomi, A. and T. Uchiyama (1998). Growth inhibiting factors in sick soil of fig orchards. *Journal of the Japanese Society for Horticultural Science* 67(1): 44-50. Osaka Prefectural Agric. Forestry Res. Cent., 442 Shakudo, Habikino-city, Osaka 583, Japan
- Katosyannos, B. I., N. A. Kouloussis, et al. (1998). Seasonal and annual occurrence of Mediterranean fruit flies (Diptera: Tephritidae) on Chios Island, Greece: Differences between two neighboring citrus orchards. *Annals of the Entomological Society of America* 91(1): 43-51. {a} Lab. Applied Zool. Parasitology, Dep. Agric., Univ. Thessaloniki, 540 06 Thessaloniki, Greece
- Kumar, V., A. Radha, et al. (1998). In vitro plant regeneration of fig (*Ficus carica* L. cv. gular) using apical buds from mature trees. *Plant Cell Reports* 17(9): 717-720. {a} Dep. Biochem., Sri Krishnadevaraya Univ., Anantapur-515 003, India
- Michailides, T. J. and D. P. Morgan (1998). Spread of endosepsis in Calimyrna fig orchards. *Phytopathology* 88(7): 637-647. {a} Dep. Plant Pathol., Univ. Calif. Davis, Kearney Agric. Cent., 9240 S. Riverbend Ave., Parlier, CA 93648, USA
- Moreno, D. A., G. Pulgar, et al. (1998). Nutritional diagnosis of fig tree leaves. *Journal of Plant Nutrition* 21(12): 2579-2588. Dep. Plant Biol., Fac. Sci., Univ. Granada, Fuentenueva Av. S/N, E-18071 Granada, Spain
- Nirenberg, H. I. and D. K. O (1998). New *Fusarium* species and combinations within the *Gibberella fujikuroi* species complex. *Mycologia* 90(3): 434-458. {a} Biol. Bundesanstalt Land Forstwirtschaft, Inst. Mikrobiol., Koenigin-Luise-Strasse 19, D-14195 Berlin, Germany
- Park, Y. S., B. W. Kim, et al. (1998). Effects of storage temperature and CO₂-enriched atmospheres on the shelf life of fig fruits during storage. *Journal of the Korean Society for Horticultural Science* 39(2): 184-188. {a} Dep. Hortic. Breeding, Mokpo Natl. Univ., Chunggae 534-729, South Korea
- Serraclara, A., F. Hawkins, et al. (1998). Hypoglycemic action of an oral fig-leaf decoction in type-I diabetic patients. *Diabetes Research and Clinical Practice* 39(1): 19-22. {a} Dep. Physiol., Fac. Med., Badajoz and Serv. Endocrinol., Univ. Hosp., 12 Octubre, Madrid, Spain
- Sforza, R., D. Clair, et al. (1998). The role of *Hyalesthes obsoletus* (Hemiptera: Cixiidae) in the occurrence of bois noir of grapevines in France. *Journal of Phytopathology Berlin* 146(11-12): 549-556. {a} Inst. Natl. Rech. Agron., Stn. Rech. Phytoplasmes, 21034 Dijon Cedex, France

Teragishi, A., Y. Kanbara, et al. (1998). Comparison between phytotron- and glasshouse-propagated fig (*Ficus carica* L.) trees grown under similar solution culture on subsequent fruiting, plant growth and fruits quality. *Journal of the Japanese Society for Horticultural Science* 67(5): 715-720. Kyoto Prefectural Yamashiro Horticultural Res. Inst., Tanabe, Tsuzuki, Kyoto 610-03, Japan

Awamura, M., D. Yahata, et al. (1997). Differences in fruit weight and soluble solids content in the juice from caprifig and fig types, and first or second crops among fig (*Ficus carica* L.) hybrid seedlings. *Journal of the Japanese Society for Horticultural Science* 66(1): 77-84. Buzen Branch, Fukuoka Agric. Res. Cent., Yukuhashi, Fukuoka 824, Japan

Baldini, E., O. Facini, et al. (1997). Leaf characteristics and optical properties of different woody species. *Trees Berlin* 12(2): 73-81. {a} Dip. Colture Arboree, Univ. Bologna, Via F. Re 6, I-40126 Bologna, Italy

Clifford, S. C., I. Kadzere, et al. (1997). Field comparisons of photosynthesis and leaf conductance in *Ziziphus mauritiana* and other fruit tree species in Zimbabwe. *Trees Berlin* 11(7): 449-454. {a} Horticulture Res. Int., Wellesbourne, Warwick CV35 9EF, UK

Gibernau, M., H. R. Buser, et al. (1997). Volatile compounds from extracts of figs of *Ficus carica*. *Phytochemistry Oxford* 46(2): 241-244. CEFE/CNRS, 1919 Route de Mende, 34293 Montpellier Cedex 5, France

Havaux, M. and F. Tardy (1997). Photoacoustically monitored thermal energy dissipation and xanthophyll cycle carotenoids in higher plant leaves. *Journal of Photochemistry and Photobiology B Biology* 40(1): 68-75. {a} Departement d'Ecophysiologie Vegetale et de Microbiologie, CEA-Cadarache, F-13108 Saint-Paul-lez-Durance, France

Mushin, T. M. and J. N. Nema (1997). Interaction between endomycorrhizae and rhizosphere fungi in soils of Iraq. *Acta Mycologica* 32(1): 41-50. Dep. Biol., Coll. Educ., Basrah Univ. Basrah, Iraq

Okubo, M. and N. Utsunomiya (1997). Effects of NaCl and temperature on growth and flower bud formation of 'Masui Dauphine' fig trees. *Environment Control in Biology* 35(3): 177-183. {a} Faculty Agriculture, Kyoto Univ., Kyoto 606, Japan

Takagaki, M., Y. Udagawa, et al. (1997). Effect of pretreatment on rooting and shoot growth of *Ficus carica* L. cuttings. *Technical Bulletin of Faculty of Horticulture Chiba University*(51): 227-230. {a} Univ. Farm, Fac. Hortic., Chiba Univ., Matsudo, Japan

Antunes, L. E. C., N. N. J. Chalfun, et al. (1996). Influence of different periods of stratification, indolebutyric acid concentration and substrate on rooting of fig cuttings. *Ciencia e Agrotecnologia* 20 (3): 307-314. {a} Dep. Agric., Universidade Federal de Lavras, C.P. 37,200-000 Lavras-MG, Brazil

Awamura, M., K. Shoda, et al. (1996). Effect of various seed parents on frequency distribution of parthenocarpy among seedling progenies of fig (*Ficus carica* L.). *Journal of the Japanese Society for*

Horticultural Science 65(1): 21-26. Buzen Banch, Fukuoka Agric. Res. Cent. Yukuhashi, Fukuoka 824, Japan

Brown, P. H. and H. Hu (1996). Phloem mobility of boron is species dependent: Evidence for phloem mobility in sorbitol-rich species. *Annals of Botany London* 77(5): 497-505. Dep. Pomology, Univ. California-Davis, Davis, CA 95616, USA

Dickson, C. (1996). Food, medicinal and other plants from the 15th century drains of Paisley Abbey, Scotland. *Vegetation History and Archaeobotany* 5(1-2): 25-31. Dep. Environ. Evol. Biol., Graham Kerr Build., Univ. Glasgow, Glasgow G12 8QQ, UK

Doster, M. A., T. J. Michailides, et al. (1996). *Aspergillus* species and mycotoxins in figs from California orchards. *Plant Disease* 80(5): 484-489. {a} Dep. Plant Pathol., Univ. Calif. Davis, Kearney Agric. Cent., 9240 S. Riverbend Ave., Parlier, CA 93648, USA

Gibernau, M., M. M. Hossaert, et al. (1996). Consequences of protecting flowers in a fig: A one-way trip for pollinators? *Journal of Biogeography* 23(4): 425-432. {a} Centre d'Ecologie Fonctionnelle et Evolutive, CNRS, BP 5051, 34033 Montpellier Cedex, France

Hernandez, R. J. A. and L. F. J. Perez (1996). New data on *Alloxantha lutea* Seidlitz, 1899 (Coleoptera: Oedemeridae). *Boletín de la Asociación Española de Entomología* 20(1-2): 255-256. {a} Pedro Antonio de Alarcón 73, 4 B., 18003 Granada, Spain

Laisk, A. and F. Loreto (1996). Determining photosynthetic parameters from leaf CO₂ exchange and chlorophyll fluorescence: Ribulose-1,5-bisphosphate carboxylase/oxygenase specificity factor, dark respiration in the light, excitation distribution between photosystems, alternative electron transport rate, and mesophyll diffusion resistance. *Plant Physiology Rockville* 110(3): 903-912. {a} Tartu Ülikooli Molekulaar-ja Rakubiolo. Inst., Riia tn. 181, Tartu EE2400, Estonia

Melanitou, M. A., D. N. Papanicolaou, et al. (1996). Sulphur dioxide binding and losses during processing and storage of dried figs. *Sciences des Aliments* 16(5): 505-513. {a} National Agricultural Res. Foundation, Inst. Technology Agricultural Products, 1, S. Venize Lou, Lycovrissi 141 23, Greece

Michailides, T. J., D. P. Morgan, et al. (1996). An old disease still a dilemma for California growers. *Plant Disease* 80(8): 828-841. {a} 9240 South Riverbend Ave., Parlier, CA 93648, USA

Noeth, M., F. Feindt, et al. (1996). Extraction of DNA and Random Amplified Polymorphic DNA (RAPD) analysis of *Ficus carica* L. and *Ficus sycomorus* L. *Mitteilungen aus dem Institut fuer Allgemeine Botanik Hamburg* 26: 153-160. Inst. Allgemeine Bot., Bot. Garten, Univ. Hamburg, Ohnhorststrasse 18, D-22609 Hamburg, Germany

Okubo, M. and N. Utsunomiya (1996). Effects of sodium chloride on growth, gas exchange and ion concentration in latex of fig (*Ficus carica* L.). *Environment Control in Biology* 34(4): 259-265. {a} Exp. Farm, Fac. Agric., Kyoto Univ., Takatsuki 569, Japan

Shiraishi, S. C., K. Kawakami, et al. (1996). Organic acid profiles in the juice of fig fruits. *Journal of the Faculty of Agriculture Kyushu University* 41(1-2): 29-33. Fruit Sci. Lab., Fac. Agric., Kyushu Univ., Fukuoka 811-23, Japan

Subbarao, K. V. and T. J. Michailides (1996). Development of phenological scales for figs and their relative susceptibilities to endosepsis and smut. *Plant Disease* 80(9): 1015-1021. {a} Dep. Plant Pathol., Univ. California Davis, c/o US Agric. Res. Station, 1636 E. Alisal St., Salinas, CA 93905, USA

Vovlas, N. and A. Larizza (1996). Relationship of *Schistonchus caprifici* (Aphelenchoididae) with fig inflorescences, the fig pollinator *Blastophaga psenes*, and its cleptoparasite *Philotrypesis caricae*. *Fundamental and Applied Nematology* 19(5): 443-448. Ist. Nematol. Agraria, Consiglio Nazionale delle Ricerche, Via Amendola 165/A, 70126 Bari, Italy

Yahata, D. and M. Awamura (1996). Varietal differences and changes in cell wall composition in ripening fig fruits. *Journal of the Japanese Society for Horticultural Science* 65(1): 41-47. Buzen Branch, Fukuoka Agric. Res. Cent., Nishiizumi, Yukuhashi, Fukuoka 824, Japan

Awamura, M., K. Shoda, et al. (1995). In Vitro Germination of Caprifig Pollen Grains using Stigmatic Exudate of a Common Fig as a Promoter. *Journal of the Japanese Society for Horticultural Science* 63(4): 739-743. Buzen Branch, Fukuoka Agric. Res. Cent., Yukuhashi, Fukuoka 824, Japan

Dechamp, C., J. C. Bessot, et al. (1995). First report of anaphylactic reaction after fig (*Ficus carica*) ingestion. *Allergy Copenhagen* 50(6): 514-516. {a} Immuno-Virol. Mol. Cellulaire, CNRS UMR 30, Fac. Med. Alexis Carrel, rue Guillaume Paradin, 69372 Lyon Cedex 08, France

Goula, M. and F. N. Ribeiro (1995). *Malacocoris chlorizans* (Panzer, 1794), new species in Portugal (Insecta, Heteroptera, Miridae). *Boletim da Sociedade Portuguesa de Entomologia*(152): 9-15. {a} Dep. Biol. Anim., Avda Diagonal, 645. E-08028 Barcelona, Spain

Khadari, B., M. Gibernau, et al. (1995). When figs wait for pollinators: The length of fig receptivity. *American Journal of Botany* 82(8): 992-999. {a} Cent. d'Ecol. Fonctionnelle et Evol., CNRS, BP 5051 34033 Montpellier Cedex, France

Khadari, B., P. Lashermes, et al. (1995). RAPD fingerprints for identification and genetic characterization of fig (*Ficus carica* L.) genotypes. *Journal of Genetics and Breeding* 49(1): 77-85. {a} Dep. Biol. Polupations, Cent. Ecologie Fonctionnelle Evolutive, CNRS 1919, route de Mende, B.P. 5051, 34033 Montpellier Cedex, France

Krajewski, D., W. Neugebauer, et al. (1995). Enantiodifferentiation of solerol isomers in dried figs and dates. *Zeitschrift fuer Lebensmittel Untersuchung und Forschung* 201(4): 378-380. {a} Lehrstuhl Lebensmittelchemie, Univ. Wuerzburg, D-97074 Wuerzburg, Germany

Mao, L., T. Ying, et al. (1995). Respiration rate, ethylene production, and cellular leakage of fig fruit following vibrational stress. *Hortscience* 30(1): 145. Dep. Food Sci., Zhejiang Agric. Univ., Hangzhou

310029, China

Massmanian, A. (1995). Severe contact dermatitis due to natural latex. *Contact Dermatitis* 32(6): 364-365. Dep. Dermatol., Hosp. Univ. La Fe, Avenguda de Campanar 21, 46009 Valencia, Spain

Oginuma, K. and H. Tobe (1995). Karyomorphology of some Moraceae and Cecropiaceae (Urticales). *Journal of Plant Research* 108(1091): 313-326. {a} Coll. Child Development, Kochi Women's Univ., 132 Ohara-cho, Kochi, 780, Japan

Subbarao, K. V. and T. J. Michailides (1995). Effects of temperature on isolates of *Fusarium moniliforme* causing fig endosepsis and *Aspergillus niger* causing smut. *Phytopathology* 85(6): 662-668. {a} Dep. Plant Pathol., Univ. Calif. Davis, c/o U.S. Agric. Res. Station, 1636 E. Alisal St., Salinas, CA 93905, USA

Summers, C. G., A. S. Newton, Jr., et al. (1995). Susceptibility of selected grape cultivars and tree fruit to silverleaf whitefly (*Bemisia argentifolii*) colonization. *Hortscience* 30(5): 1040-1042. {a} Univ. Calif., 9240 S. Riverbend Ave., Parlierr, CA 93648, USA

Vicidomini, S. (1995). Biology of *Xylocopa (Xylocopa) violacea* (L., 1758) (Hymenoptera: Apidae): Nest morphology. *Atti della Societa Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano* 136(2): 95-107. {a} Univ. Federico II, Dip. di Zool., via Mezzocannone 8, 80134 Napoli, Italy

Yoshioka, K. (1995). Changes in saccharides and enzyme activities related to carbohydrate metabolism during growth and development of fig fruit (*Ficus carica* L.) on the tree. *Journal of the Japanese Society for Food Science and Technology* 42(4): 219-224. Dep. Clin. Nutr. Sci., Osaka Prefectural Junior Coll. Health Sci, 3-7-30 Habikino, Habikino-shi 583, Japan

Bacic, T. and M. Ruscic (1994). Acidification and buffer capacity of bark in five deciduous plants from the region of Split (Croatia). *Acta Biologica Cracoviensia Series Botanica* 36: 51-56. {a} Fac. Educ., Univ. J. J. Strossmayer, L. Jaegera 9, 54000 Osijek, Croatia

Brown, P. H. (1994). Seasonal variations in fig (*Ficus carica* L.) leaf nutrient concentrations. *Hortscience* 29(8): 871-873. Dep. Pomol., Univ. Calif., Davis, CA 95616-8683, USA

Canard, M. (1994). A cottony scale *Pulvinaria horii* Kuwana, 1902 (Homoptera: Coccidae) new to the fauna of continental Europe. *Biologia Gallo Hellenica* 21(1): 35-40. 47 Chemin Flou-de-Rious, F-31400 Toulouse, France

Golombek, S. D., B. C. Hecht, et al. (1994). Ultrastructure of the endodermis of fig root tips in response to salinity. *Angewandte Botanik* 68(3-4): 79-82. {a} Dep. Agronomy, International Crops Res. Inst., Semi-Arid Tropics, Patancheru PO, Andhra Pradesh 502324, India

Hossaert, M. M., M. Gibernau, et al. (1994). Chemosensory attraction of fig wasps to substances produced by receptive figs. *Entomologia Experimentalis et Applicata* 70(2): 185-191. {a} CEFE, CNRS,

Route de Mende, BP 5051, 34033 Montpellier Cedex, France

Lisci, M. and E. Pacini (1994). Germination ecology of drupelets of the fig (*Ficus carica* L.). *Botanical Journal of the Linnean Society* 114(2): 133-146. {a} Dep. Plant Biol., Univ. Florence, Via G. La Pira 4, I-50121 Florence, Italy

Macfall, J. S. and G. A. Johnson (1994). The architecture of plant vasculature and transport as seen with magnetic resonance microscopy. *Canadian Journal of Botany* 72(11): 1561-1573. {a} Sch. Environment, Dep. Radiol., Duke Univ., Durham, NC 27710, USA

Michailides, T. J. and D. P. Morgan (1994). Dynamics of *Blastophaga psenes* populations, availability of caprifigs, and fig endosepsis caused by *Fusarium moniliforme*. *Phytopathology* 84(11): 1254-1263. {a} Dep. Plant Pathol., Univ. Calif., Davis, CA, USA

Michailides, T. J., D. P. Morgan, et al. (1994). Comparison of three methods for determining fig endosepsis caused by *Fusarium moniliforme* and other molds in caprifigs and calimyrna figs. *Plant Disease* 78(1): 44-50. Dep. Plant Pathology, Univ. Calif., Davis, CA, USA

Siddiqui, M. B. and W. Husain (1994). Medicinal plants of wide use in India with special reference to Sitapur district (Uttar Pradesh). *Fitoterapia* 65(1): 3-6. Dep. Botany, Aligarh Muslim Univ., Aligarh 202002, India

Golombek, S. D. and P. Luedders (1993). Effects of short-term salinity on leaf gas exchange of the fig (*Ficus carica* L.). *Plant and Soil* 148(1): 21-27. Inst. Crop Sci., Technical Univ. Berlin, Albrecht-Thaer Weg 3, D-1000 Berlin 33, Germany

Honma, Y., K. Minowa, et al. (1993). Remarkable strandings of Pacific white sided dolphins, *Lagenorhynchus obliquidens* Gill, 1865, and other cetaceans on the coast of Niigata District, Sea of Japan, during severe winter conditions, 1992-1993. *Bulletin of the Biogeographical Society of Japan* 48(1): 81-86. {a} Fac. Science, Niigata Univ., Ikarashi 2, Niigata 950-21, Japan

Mann, J. S. and A. S. Sohi (1993). A new species of genus *Goifa dworakowska* (Homoptera: Cicadellidae) from northern India. *Journal of Insect Science* 6(1): 97-98. Dep. Entomol., Punjab Agric. Univ., Ludhiana-141 004, India

Mathooko, F. M., T. Sotokawa, et al. (1993). Retention of freshness in fig fruit by CO₂-enriched atmosphere treatment or modified atmosphere packaging under ambient temperature. *Journal of the Japanese Society for Horticultural Science* 62(3): 661-667. {a} Graduate Sch. Natural Sci. and Technol., Okayama Univ., Tsushima, Okayama 700, Japan

Parisi, L., Y. Lespinasse, et al. (1993). A new race of *Venturia inaequalis* virulent to apples with resistance due to the Vf gene. *Phytopathology* 83(5): 533-537. {a} INRA, Station Pathologie, Vegetale Phytobacteriologie, Rue Georges Morel, B.P. 57, 49071 Beaucouze Cedex, France

- Subbarao, K. V. and T. J. Michailides (1993). Virulence of *Fusarium* species causing fig endosepsis in cultivated and wild caprifigs. *Phytopathology* 83(5): 527-533. {a} Dep. Plant Pathol., Univ. Calif. Davis, U.S. Agric. Res. Station, 1636 E. Alisal Street, Salina 93905, USA
- Bosch, M., V. Pedrocchi, et al. (1992). The heronry of the Medes Islands (NE Spain). *Miscellanea Zoologica Barcelona* 16: 249-253. Dep. Biol. Animal, Fac. de Biol., Univ. de Barcelona, Avgda. Diagonal 645, 08028 Barcelona, Spain
- Dickson, J. H. (1992). Some recent additions to the Quaternary flora of Scotland and their phytogeographical, palaeoclimatic and ethnobotanical significance: A review. *Acta Botanica Fennica* 144: 51-57.
- Hornyts, k. I. P. and H. H. Martyn (1992). Properties of the microscopic structure of *Ficus* species as criteria of their selection for interior decorating. *Ukrayins'kyi Botanichnyi Zhurnal* 49(1): 34-39. {a} Donetsk Bot. Gard., Acad. Sci. Ukr., Donetsk, Ukraine
- Karlsson, T. (1992). Floristical notes. *Svensk Botanisk Tidskrift* 86(2): 93-94.
- Kim, S. S., C. H. Lee, et al. (1992). Chemical components in the two cultivars of Korean figs (*Ficus carica* L.). *Journal Of The Korean Agricultural Chemical Society* 35(1): 51-54.
- Kimura, N., M. Sakakibara, et al. (1992). Lighting for coloring and quality in fig fruit in maturing period. *Research Bulletin of the Aichi ken Agricultural Research Center*(24): 195-204. Aichi-ken Agric. Res. Cent., Nagakute, Aichi 480-11, Japan
- Kucherenko, V. P., V. I. Severin, et al. (1992). Prediction of plant winter resistance from peroxidase activity indices. *Fiziologiya i Biokhimiya Kul'turnykh Rastenii* 24(2): 179-182. A.V. Fomin Bot. Gard., T. G. Shevchenko Kiev State Univ., Kiev, Ukraine
- Luedders, P. and V. Kaminski (1992). Influence of soil salinity on water consumption and transpiration of two fig cultivars (*Ficus carica* L.) in different air humidity. *Angewandte Botanik* 66(1-2): 31-35.
- McSorley, R. (1992). Nematological problems in tropical and subtropical fruit tree crops. *Nematropica* 22(1): 103-116.
- Smillie, R. M. (1992). Calvin cycle activity in fruit and the effect of heat stress. *Scientia Horticulturae* 51 (1-2): 83-95.
- Subbarao, K. V. and T. J. Michailides (1992). A re-evaluation of *Fusarium moniliforme* var. *fici*, the causal agent of fig endosepsis. *Mycological Research* 96(9): 766-768.
- Van, D. V. M. (1992). Botanical evidence for Garamantian agriculture in Fezzan, southern Libya. *Review of Palaeobotany and Palynology* 73(1-4): 315-327. Sch. Archaeol. Studies, Univ. Leicester, University Rd., Leicester LE1 7RH, UK

- Vovlas, N., R. N. Inserra, et al. (1992). *Schistonchus caprifici* parasitizing caprifig (*Ficus carica sylvestris*) florets and the relationship with its fig wasp (*Blastophaga psenes*) vector. *Nematologica* 38 (2): 215-226.
- Winter, K. and M. Lesch (1992). Diurnal changes in chlorophyll a fluorescence and carotenoid composition in *Opuntia ficus indica*, a CAM plant, and in three C-3 species in Portugal during summer. *Oecologia Heidelberg* 91(4): 505-510. {a} Smithsonian Tropical Res. Inst., Box 2072, Balboa, Panama
- Zervakis, G., D. Dimou, et al. (1992). First record of the natural occurrence in Europe of the basidiomycete *Pleurotus cystidiosus* on a new host. *Mycological Research* 96(10): 874-876. Lab. Gen. Agric. Microbiol., Agric. Univ. Athens, Oera Odos 75, 11855 Athens, Greece
- Abdel, R. M. S. and D. S. El (1991). Functional adaptations of fig trees *Ficus carica*, L.) in agroecosystems of the western Mediterranean desert of Egypt. *Qatar University Science Bulletin* 11: 183-199. {a} Dep. Botany, Fac. Sci., Univ. Qatar, Doha, Qatar
- Burla, H. and G. Bachli (1991). Notes on rearing drosophilid species from both cultivated and natural fruits. *Mitteilungen Der Schweizerischen Entomologischen Gesellschaft* 64(1-2): 45-53.
- Flach, B. M. T. and B. M. Eller (1991). Solar energy input to plant surfaces: III. Influence of leaf curvature and self-shading. *Botanica Helvetica* 101(1): 121-134.
- Sultan, S. A. and H. Ferris (1991). The effect of soil moisture and soil particle size on the survival and population increase of *Xiphinema index*. *Revue De Nematologie* 14(3): 345-352.
- Carballo, G. B. M. (1990). Use of the Calabacita variety of fig (*Ficus carica*) in the production of jam. *Alimentaria* 27(213): 63-64.
- Nii, N. and T. Kuroiwa (1990). Changes of chloroplast ultrastructure and plastid nucleoids during greening under light in etiolated fig leaves (*Ficus carica*). *Journal Of The Japanese Society For Horticultural Science* 59(2): 333-340.
- Pradhan, K. C. and S. N. Das (1990). Further observations on nematodes associated with high altitude plants, forest and medicinal plants of Phulbani District (Orissa) (India). *Indian Forester* 116(2): 163-167.
- Sohi, A. S. and J. S. Mann (1990). A new species of genus *Daluana* Ramakrishnan (Cicadellidae: Typhlocybae: Empoascini) infesting fig at Rishikesh (Uttar Pradesh) India. *Entomon* 15(3-4): 267-270.
- Tsantili, E. (1990). Changes during development of Tsapela fig fruits. *Scientia Horticulturae* 44(3-4): 227-234.

- Bacic, T., S. Gligorijevic, et al. (1989). Effect of air pollution on ascorbic acid content in fruits of some Mediterranean trees. *Acta Biologiae Et Medicinae Experimentalis* 14(2): 151-156.
- Bollet, C., C. Gulian, et al. (1989). Isolation of *Serratia rubidaea* (Stapp) from fresh and spoiled coconuts. *Tropical Agriculture* 66(4): 342-344.
- Cormier, F., C. Charest, et al. (1989). Partial purification and properties of proteases from fig (*Ficus carica*) callus cultures. *Biotechnology Letters* 11(11): 797-802.
- Kaminski, V. and P. Luedders (1989). Influence of sodium chloride on vegetative growth of *Ficus carica* L. at different relative atmospheric humidity. *Gartenbauwissenschaft* 54(2): 58-61.
- Kim, S. S. and J. H. Lee (1989). A taxonomic study of some species in genus *Ficus* (Moraceae). *Korean Journal Of Plant Taxonomy* 19(2): 123-138.
- Kulieva, A. A. (1989). Data on the mycobiota of wild fruit plants in the Azerbaijan SSR (USSR). *Izvestiya Akademii Nauk Azerbaidzhanskoi Ssr Seriya Biologicheskikh Nauk*(1): 122-126.
- Maruyama, S., S. Miyoshi, et al. (1989). Angiotensin I-converting enzyme inhibitors derived from *Ficus carica*. *Agricultural And Biological Chemistry* 53(10): 2763-2768.
- Rapisarda, C. (1989). On the presence in Sicily (Italy) of the green fig psyllid (*Homotoma viridis* Klimaszewski) and description of its fifth instar nymph (Homoptera Psylloidea). *Bollettino Della Societa Entomologica Italiana* 121(1): 13-18.
- Shorey, H. H., L. Ferguson, et al. (1989). Solar heating reduces insect infestations in ripening and drying figs. *Hortscience* 24(3): 443-445.
- Alwan, A. H. and B. Z. A. F. Al (1988). Effects of milk latex of fig (*Ficus carica*) on tritiated benzo(a) pyrene binding to rat liver microsomal protein. *International Journal Of Crude Drug Research* 26(4): 209-213.
- De, L. E. (1988). New eriophyid mites (Acari: Eriophyoidea) from Italy: I. *Entomologica* 23: 13-46.
- Ito, M. F. and F. A. Bergamin (1988). Inoculation methods of *Ceratocystis fimbriata* Ell. and Halst. on *Acacia decurrens*, *Crotalaria juncea*, *Ficus carica*, *Mangifera indica* and pathogenicity in different hosts. *Summa Phytopathologica* 14(1-2): 117-135.
- Kjellberg, F., B. Doumesche, et al. (1988). Longevity of a fig wasp (*Blastophaga psenes*). *Proceedings Of The Koninklijke Nederlandse Akademie Van Wetenschappen Series C Biological And Medical Sciences* 91(2): 117-122.
- Puri, S. and R. Nagpal (1988). Effect of auxins on air-layers of some agro-forestry species. *Indian Journal Of Forestry* 11(1): 28-32.

- Wyss, U., W. M. Robertson, et al. (1988). Esophageal bulb function of *Xiphinema index* and associated root cell responses, assessed by video-enhanced contrast light microscopy. *Revue De Nematologie* 11 (2): 253-262.
- Jensen, H. A. (1987). Macrofossils and their contribution to the history of the spermatophyte flora in Southern Scandinavia from 13000 BP to 1536 AD. *Kongelige Danske Videnskabernes Selskab Biologiske Skrifter* 29: 1-68.
- Kjellberg, F., P. H. Gouyon, et al. (1987). The stability of the symbiosis between dioecious figs and their pollinators: A study of *Ficus carica* L. and *Blastophaga psenes* L. *Evolution* 41(4): 693-704.
- Nassar, A. H. and H. J. Newbury (1987). Ficin production by callus cultures of *Ficus carica*. *Journal Of Plant Physiology* 131(3-4): 171-180.
- Ribeiro, I. J. A., M. F. Ito, et al. (1987). *Cassia renigera* Wall.: A new host of *Ceratocystis fimbriata* Ell. and Halst. *Bragantia* 46(2): 417-424.
- Valizadeh, M., G. Valdeyron, et al. (1987). Gene flow in the common fig, *Ficus carica*: pollen dispersal in a dense stand. *Acta Oecologica Oecologia Plantarum* 8(2): 143-154.
- Brighenti, E., F. B. G. Santos, et al. (1986). Influence of the intensity of pruning on fructification in the fig tree (*Ficus carica* L.) on green fig production, Roxo de Valhinas cultivar. *Agros* 21(1-4): 67-75.
- Golubev, V. N., L. N. Pilipenko, et al. (1986). Fractionation and composition of carbohydrates from *Ficus carica*. *Khimiya Prirodnikh Soedinenii*(6): 673-677.
- Kolesnik, A. A., T. A. Kakhniashvili, et al. (1986). Lipids from *Ficus carica* fruits. *Khimiya Prirodnikh Soedinenii*(4): 423-427.
- McKenzie, E. H. C. (1986). New plant disease record in New Zealand: Fig rust (*Cerotelium fici*) on *Ficus carica*. *New Zealand Journal Of Agricultural Research* 29(4): 707-710.
- Nagpal, R. and S. Puri (1986). Effect of auxins on air-layers of some agro-forestry species. *Indian Journal Of Forestry* 9(3): 232-236.
- O'Toole, J. C. and J. G. Real (1986). Estimation of aerodynamic and crop resistance from canopy temperature. *Agronomy Journal* 78(2): 305-310.
- Pisarnitskii, A. F., I. A. Egorov, et al. (1986). Scent-forming components of fig (*Ficus carica*) fruits. *Prikladnaya Biokhimiya I Mikrobiologiya* 22(2): 281-285.
- Brown, D. J. F. and M. I. Coiro (1985). The reproductive capacity and longevity of *Xiphinema index* (Nematoda: Dorylaimida) from three populations on selected host plants. *Revue De Nematologie* 8(2):

171-174.

Credaro, V. and A. Pirola (1985). News about frost damages for cultivated plants in Lombardy (Italy), 1985. *Atti Dell'Istituto Botanico Della Universita E Laboratorio Crittogamico Pavia* 4(Series 7): 27-54.

Leporatti, M. L., E. Posocco, et al. (1985). Some new therapeutic uses of several medicinal plants in the Province of Terni (Umbria, Central Italy). *Journal Of Ethnopharmacology* 14(1): 65-68.

Nagpal, R. and R. N. Sehgal (1985). Propagation of some agro-forestry species by air-layering. *Indian Journal Of Forestry* 8(3): 161-165.

Rokba, A. M., A. H. Shahein, et al. (1985). Fruit trees collection farm in King Mariut (Egypt) 1. Agrobiological characteristics and evaluation of some fig (*Ficus carica* Risso) cultivars. *Egyptian Journal Of Horticulture* 12(2): 103-114.

Siewek, F., K. Herrmann, et al. (1985). Isomeric di-C-glycosylflavones in fig (*Ficus carica*). *Zeitschrift Fuer Naturforschung Section C Biosciences* 40(1-2): 8-12.

Coiro, M. I. and D. J. F. Brown (1984). The status of some plants as hosts for 4 populations of *Xiphinema index* (Nematoda: Dorylaimida). *Revue De Nematologie* 7(3): 283-286.

Hatzinikolis, E. N. (1984). Description of *Panonychus caricae*, new species on fig trees in Greece (Acari: Tetranychidae). *Entomologia Hellenica* 2(2): 55-62.

Kadam, B. A. and V. K. Patil (1984). Effects of nitrogen, phosphorus and potassium on fig (*Ficus carica*). *Journal Of Maharashtra Agricultural Universities* 9(1): 22-24.

Patil, P. K. and V. K. Patil (1983). Salinity tolerance in fig (*Ficus carica*). P.K.: 12-14.

Pedrotti, E. L., I. Manica, et al. (1983). Irrigation levels, plant growth and nutrient concentration in the leaves of fig (*Ficus carica*) (cultivar) Roxo de Valinhos. *Agronomia Sulriograndense* 19(1): 103-116.

Beck, N. G. and E. M. Lord (1913). Breeding system in *Ficus carica*, the common fig: II. Pollination events. *American Journal Of Botany* 75(12): 1913-1922.

Beck, N. G. and E. M. Lord (1904). Breeding system in *Ficus carica*, the common fig: I. Floral diversity. *American Journal Of Botany* 75(12): 1904-1912.

Colelli, G., F. G. Mitchell, et al. (1993). Extension of postharvest life of Mission figs by carbon dioxide enriched atmospheres. *Hortscience* 26(9): 1193-1195.

Bernabo, E. Brief notes on the biology of *Liparthrum mori* (Aube) and on the xylophagous coleopterous fauna of fig (Coleoptera). *Memorie della Societa Entomologica Italiana* 71(2): 543-549.
Via A. Crocco 3G/10, 16122 Genova, Italy

Siddiqui, M. A., A. Haseeb, et al. Control of plant-parasitic nematodes by soil amendments with latex bearing plants. Indian Journal of Nematology 22(1): 25-28. {a} Dep. Bot., Aligarh Muslim Univ., Aligarh 202 002, India

Stino, R. G. and F. E. El Fruiting of Conadria fig trees as affected by type of pruning and hydrogen cyanamide application. Egyptian Journal of Horticulture 22(2): 193-211. {a} Fac. Agric., Cairo Univ., Cairo, Egypt

Index of botanical names:

[\[A\]](#) [\[B\]](#) [\[C\]](#) [\[D\]](#) [\[E\]](#) [\[F\]](#) [\[G\]](#) [\[H\]](#) [\[I\]](#) [\[J\]](#) [\[K\]](#) [\[L\]](#) [\[M\]](#)
[\[N\]](#) [\[O\]](#) [\[P\]](#) [\[Q\]](#) [\[R\]](#) [\[S\]](#) [\[T\]](#) [\[U\]](#) [\[V\]](#) [\[W\]](#) [\[X\]](#) [\[Y\]](#) [\[Z\]](#)

Index of common names:

[\[A\]](#) [\[B\]](#) [\[C\]](#) [\[D\]](#) [\[E\]](#) [\[F\]](#) [\[G\]](#) [\[H\]](#) [\[I\]](#) [\[J\]](#) [\[K\]](#) [\[L\]](#) [\[M\]](#)
[\[N\]](#) [\[O\]](#) [\[P\]](#) [\[Q\]](#) [\[R\]](#) [\[S\]](#) [\[T\]](#) [\[U\]](#) [\[V\]](#) [\[W\]](#) [\[X\]](#) [\[Y\]](#) [\[Z\]](#)

All information is included in good faith but The University of Queensland does not warrant or guarantee the accuracy of any information on these pages, nor does it accept responsibilities for any loss arising from the use of this information. Views and opinions do not represent those of the University of Queensland.

Contact: Dr Rob Fletcher, School of Agriculture and Horticulture, The University of Queensland, Gatton 4343; Telephone: 07 5460 1311 or 07 5460 1301; Facsimile: 07 5460 1112; International facsimile: 61 7 5460 1112; Email: r.fletcher@mailbox.uq.edu.au

[\[New Crops Home Page\]](#) [\[New Crops Program\]](#) [\[Australian New Crops
\(Periodical\)\]](#) [\[New Crops Publications\]](#) [\[Order Form\]](#) [\[People\]](#) [\[Crop Profiles\]](#)
[\[Other Resources\]](#)

Latest update 30 January 2001 by: [RF](#)