

## Herbaceous cuttings of fig

**Fig cuttings generally take place in late winter, with branches in dormancy. But a fig cuttings in early summer in full fruiting period, is quite feasible. A variety to save that can not wait for winter? A fig fruit delicious encountered strolling around? Here's how.**

The grass cutting is done in June-July.

It carries out the harvesting of herbaceous branches (shoots year green) of the variety to multiply:



It removes the leaves and cut into pieces 4 to 5 eyes. It should be noted that one can leave a terminal bud with a small leaf.

### Categories

- [General principles and methodology](#)
- [Glossary](#)
- [Multiplication techniques](#)
- [Grafting of fruit](#)
- [Grafting of ornamental](#)
- [Experiments](#)
- [Tools and Accessories](#)



Fill a pot deep enough, genre pot rose, a substrate (here the pure universal soil) that water copiously.



Pique on one by one the branches, by burying it at least half. Using Rooting hormone is not necessary.



It fills the pot and the branches in spacing of about 4 cm.



It covers all of a sealed transparent bag, which builds on the base with a rubber band or a string. We realize this is called a cutting to stew.



We store the pot in a place that will be permanently in the shade (under a tree ...).

We can discover the cuttings after a minimum of six weeks. (the best way to enjoy is the training of herbaceous stems from the cuttings, which is not the case with the photograph below, discovered too early).



They keep in the shade.

In winter, when leaves are falling, on deposit batch of cuttings for planting in individual pots.

P.-S.

Last year, I got 100% success with this method

A variant of herbaceous cuttings of fig:

<http://www.galgoni.com/FRA/Reproduccio/enverd.htm>

 Category: Techniques of multiplication

 Author: patrice